

In the Patients section of the abstract “irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria” (1), the two \geq symbols should be \leq symbols.

Reference

1. Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria [Abstract]. ACP J Club. 2002 May-Jun;136:82. Abstract of: Parving H-H, Lehnert H, Bröchner-Mortensen J, et al., for the Irbesartan in Patients with Type 2 Diabetes and Microalbuminuria Study Group. The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. N Engl J Med. 2001 Sep 20;345:870-8.

In the Intervention section of the abstract “Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure” (1), the two \geq symbols should be \leq symbols.

Reference

1. Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure [Abstract]. ACP J Club. 2002 May-Jun;136:83. Abstract of: Lewis EJ, Hunsicker LG, Clarke WR, et al., for the Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. N Engl J Med. 2001 Sep 20;345:851-60.

In the fourth column, last row of the Table of the abstract “overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes” (1), there should be a || symbol after the 10%.

Reference

1. Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes [Abstract]. ACP J Club. 2002 May-Jun;136:101. Abstract of: Gerber RA, Cappelleri JC, Kourides IA, Gelfand RA. Treatment satisfaction with inhaled insulin in patients with type 1 diabetes. A randomized controlled trial. Diabetes Care. 2001 Sep;24:1556-9.

In both rows of the first column, of the Table of the abstract “Montelukast moderately decreased asthma symptoms in children with persistent asthma” (1), the word system should be symptom.

Reference

1. Montelukast moderately decreased asthma symptoms in children with persistent asthma [Abstract]. ACP J Club. 2002 May-Jun;136:102. Abstract of: Knorr B, Franchi LM, Bisgaard H, et al. Montelukast, a leukotriene receptor antagonist, for the treatment of persistent asthma in children aged 2 to 5 years. Pediatrics. 2001 Sep;108:e48.

In the last row of the first column, of the Table of the abstract “1 of 2 quality-improvement interventions for depression in managed care was more effective but more costly than usual care” (1), there should be a || symbol after the word *costs*.

Reference

1. 1 of 2 quality-improvement interventions for depression in managed care was more effective but more costly than usual care [Abstract]. ACP J Club. 2002 May-Jun;136:116. Abstract of: Schoenbaum M, Unützer J, Sherbourne C, et al. Cost-effectiveness of practice-initiated quality improvement for depression. Results of a randomized controlled trial. JAMA. 2001 Sep 19;286:1325-30.